

WHY IS NATURE RESTORATION CRITICAL FOR CLIMATE MITIGATION IN THE EU?

Publication date:
December 2022

Authors:
Evelyn Underwood, Gabrielle Aubert (IEEP)



The EU has set targets to reduce its greenhouse gas emissions by at least 55% by 2030 and to reach climate neutrality by 2050 to keep global temperature increase below 1.5°C. Europe's land is vital to achieving this goal. The land use and land use change and forestry (LULUCF) sector will contribute by saving and storing an additional 310 to 400 MtCO₂eq by 2030.

It will be impossible for the EU to meet its climate neutrality goal by 2050 and the LULUCF land sink target without a significant scaling up of nature restoration.

Nature restoration is central to our efforts to mitigate climate change as it reduces and avoids emissions from land, enhances the capacity of ecosystems to capture and sequester carbon in natural sinks, and can prevent future emissions by increasing ecosystem resilience.

WHAT CAN THE NRL ACHIEVE?

- Restoring **peatlands, agroecosystems and forests** holds great potential to safeguard carbon stocks and increase sequestration.
- Theoretically, restoring 90% of terrestrial Annex I habitats which are in not good or unknown condition to good condition could lead to a total carbon stock between 2,858 and 9,210 million tC in the EU and sequester around 286 MtCO₂eq/year [1]. While the time needed to achieve these rates would largely exceed the 2030 deadline, these figures illustrate the theoretical magnitude of the carbon storage and sequestration benefits which could be achieved by Article 4 of the nature restoration law.
- Events such as forest fires, floods and droughts will become more frequent and severe with climate change over this decade. These events release large amounts of stored carbon into the atmosphere. Restoration can increase the resilience of habitats to climate events and reduce their impact on carbon stocks.

WHAT WOULD HAPPEN WITH NO RESTORATION?

- If nature restoration is not scaled up, the EEA projects a decrease in the overall level of net removals for the period 2020-2040, with an average of 200 MtCO₂e removed each year - compared to the historic average of 300 MtCO₂e for the period 1990-2019 [2]. This would mean the EU is very likely to fail to achieve carbon neutrality by 2050.

Why is nature restoration critical for climate mitigation in the EU?

Peatlands


Peatlands cover around 3% of the EU-27 agricultural area – yet emit 25% of the EU's annual emissions from agricultural land [3].

WHAT CAN THE NRL ACHIEVE?

- The restoration of peatland and wetland Annex I habitats under Article 4 could achieve additional net GHG mitigation benefits between 7.8 and 22.8 million tCO₂eq/year to 2030 and between 26.7 and 62.9 MtCO₂eq/year to 2050 [4]. The technical mitigation potential of rewetting and restoring all EU peatlands is much larger – it is estimated at about 185 MtCO₂eq/year, on average for the 2020-2050 period [5].
- Rewetting 35% of the total area of agriculturally used peatlands in the EU under Article 9.4 will **reduce their total emissions by 25% (around 45 Mt CO₂eq)** [6].

WHAT WOULD HAPPEN WITH NO RESTORATION?

The IPCC estimate that if no rewetting is done, emissions from peatland would take up 12-42% of the global emission budget needed to keep warming below 1.5-2°C [7].



Keava raba, Ohekatku, Rapla County, Estonia, Photo by Single.Earth

Agroecosystems

Restoring agroecosystems can re-establish their natural carbon cycling and storage capacities - thereby achieving climate mitigation benefits through avoiding some current emissions and enhancing carbon sinks.

WHAT CAN THE NRL ACHIEVE?

- Increasing soil organic carbon stocks on arable land in the EU under Article 9.2(b) could achieve **an additional sequestration of 50.48 million tCO₂eq/year** [8].
- The economically feasible carbon sequestration potential of EU27 agricultural soils is at least **20 million tCO₂eq/year**, whilst the maximum technically achievable carbon sequestration potential could be 200 million tCO₂eq/year [9].

WHAT WOULD HAPPEN WITH NO RESTORATION?

Under a business-as-usual scenario, soil organic carbon (SOC) is predicted to increase between 12.8 and 13.9-14.1 Gt in cropland by 2050, and from 6.7 to 8.9-9.4 Gt in pastures [5]. This is much less than what can be achieved through restoration.


Why is nature restoration critical for climate mitigation in the EU?

Forests

Forests are the largest carbon sink in the EU, sequestering around one tenth of Europe's gross CO₂ emissions in 2020 [10]. They have the highest sequestration rate of any terrestrial habitat [2]. Their natural capacities, however, are decreasing, and forests have started to become net sources of emissions in many EU member states, due to increased harvesting, climate change, forest fires, and pest damage [11-12]. The nature restoration law proposal also requires no further degradation of existing forest habitats important for biodiversity (Annex I habitats), which includes old growth forests. **Restoring, re-establishing, and adapting forest habitats to climate change can significantly contribute to climate mitigation by increasing the forest carbon sink and avoiding emissions.**

WHAT CAN THE NRL ACHIEVE?

Restoring forests and making forests more resilient to climate change is estimated to have an **additional mitigation potential in the order of 90 to 180 million tCO₂/year by 2040** [13]. Recent studies estimate significantly larger **potentials of up to 440 million tCO₂eq/year**, which could double the climate mitigation potential of forests by 2050 [4].



Eibsee, Grainau, Germany, Photo by Daniel Seßler

WHAT WOULD HAPPEN WITH NO RESTORATION?

Without restoration, forests' carbon storage potential may decline by 180 million tCO₂eq annually from 2021-2030, reducing the sink by more than 50% [14]. Many forests are already becoming net sources of emissions and this risk will increase as climate change increases the risks of forest disasters and as the intensity of forest management increases.

Not all nature restoration will increase or safeguard carbon – sometimes saving biodiversity means losing carbon but increasing long-term habitat resilience.

- On heaths, scrub habitats, and grasslands, actions that increase carbon and biomass – planting trees or adding nutrients or letting the habitats get overgrown - can destroy their biodiversity value.
- Restoration actions may remove carbon by cutting down trees planted on peat bogs or heaths, clearing overgrowth on abandoned grasslands, or cutting away over-fertile soil and vegetation.
- Restoration can prevent or minimize fire risk in these habitats, which can avoid significant emissions in future.

Why is nature restoration critical for climate mitigation in the EU?

Peatland restoration reduces emissions and global warming potential of restored sites

The LIFE PEAT RESTORE project contributed to rewetting degraded peatlands to restore their function as carbon sinks in five EU countries around the Baltic Sea (Germany, Poland, Lithuania, Latvia, and Estonia) on an area of 5,300 hectares. The preliminary results in 2021 showed that the peatland management and restoration actions support CO₂ uptake or reduction of CO₂ emissions and has shown first steps of recovery towards functional peatland ecosystems.

It is estimated that around **30% of the global warming potential of all sites was reduced**, amounting to 14,500 tons CO₂eq/year [15].



Viru bog, Kolga, Harju County, Estonia, Photo by Jaanus Jagomägi



Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or CINEA. Neither the European Union nor CINEA can be held responsible for them.

These policy briefs were written by IEEP and Ecologic Institute in the Think Sustainable Europe Network in response to the proposed EU nature restoration law.



Why is nature restoration critical for climate mitigation in the EU?

References

1. Kopsieker, L., G. Costa Domingo, and E. Underwood, Climate mitigation potential of large-scale restoration in Europe. Analysis of the climate mitigation potential of restoring habitats listed in Annex I of the Habitats Directive. 2021, Institute for European Environmental Policy.
2. EEA, Carbon stocks and sequestration in terrestrial and marine ecosystems: a lever for nature restoration? 2022, European Environmental Agency.
3. Greifswald Mire Centre, Peatlands in the EU. Common agricultural policy (CAP) after 2020. 2020, Greifswald Mire Centre: Germany.
4. European Commission, IMPACT ASSESSMENT REPORT Accompanying the document Proposal for a Regulation of the European Parliament and the Council amending Regulations (EU) 2018/841 as regards the scope, simplifying the compliance rules, setting out the targets of the Member States for 2030 and committing to the collective achievement of climate neutrality by 2035 in the land use, forestry and agriculture sector, and (EU) 2018/1999 as regards improvement in monitoring, reporting, tracking of progress and review, in COMMISSION STAFF WORKING DOCUMENT. 2021, European Commission: Brussels.
5. Andrés, P., et al., Agricultural potential in carbon sequestration: Humus content of land used for agriculture and CO₂ storage. 2022, European Parliament Policy Department for Structural and Cohesion Policies: Brussels.
6. Greifswald Mire Centre and Wetlands International, Higher ambition for peatlands in the EU Nature Restoration Law proposal. 2022, Greifswald Mire Centre and Wetlands International Europe.
7. Leifeld, J., C. Wüst-Galley, and S. Page, Intact and managed peatland soils as a source and sink of GHGs from 1850 to 2100. *Nature Climate Change*, 2019. 9: p. 1-3.
8. Lugato, E., et al., Potential carbon sequestration of European arable soils estimated by modelling a comprehensive set of management practices. *Global Change Biology*, 2014. 20(11): p. 3557-3567.
9. Smith, P., Agricultural greenhouse gas mitigation potential globally, in Europe and in the UK: what have we learnt in the last 20 years? *Global Change Biology*, 2012. 18(1): p. 35-43.
10. Grassi, G., et al., On the realistic contribution of European forests to reach climate objectives. *Carbon Balance and Management*, 2019. 14(1): p. 8.
11. Booth, M.S., Burning up the carbon sink: How the EU's forest biomass policy undermines climate mitigation, and how it can be reformed. 2022, Partnership for Policy Integrity.
12. Statistics Finland, Greenhouse gas emissions in 2021 remained on level with the previous year, the land use sector a net source of emissions for the first time. 2022, Statistics Finland: Helsinki, Finland.
13. Nabuurs, G.-J., et al., By 2050 the mitigation effects of EU forests could nearly double through climate smart forestry. *Forests*, 2017. 8(12): p. 484.
14. Seidl, R., et al., Increasing forest disturbances in Europe and their impact on carbon storage. *Nature Climate Change*, 2014. 4(9): p. 806-810.
15. Jurema, L., et al., Layman report LIFE Peat Restore, Restoring peatlands for climate. 2022.