

# Comparison of the Efficacy of Five Front-of-Pack Nutrition Labels in Helping the Brazilian Consumer Make a Healthier Choice

Wendy A.M. Blom<sup>1\*</sup>, Nathalie C. Goenee<sup>2</sup>, Lucia Juliano<sup>3</sup>, Els M. de Groene<sup>1</sup> and Fernanda de Oliveira Martins<sup>4</sup>

<sup>1</sup>Unilever Foods Innovation Centre Wageningen, Netherlands.

<sup>2</sup>Unilever R&D Vlaardingen, Netherlands.

<sup>3</sup>Harris Interactive UK Ltd, Netherlands.

<sup>4</sup>Unilever Brazil.

## \*Correspondence:

Wendy A.M. Blom, Unilever Foods Innovation Centre Wageningen, Netherlands, E-mail: wendy.blom@unilever.com.

**Received:** 02 August 2021; **Accepted:** 28 August 2021

**Citation:** Blom WAM, Goenee NC, Juliano L, et al. Comparison of the Efficacy of Five Front-of-Pack Nutrition Labels in Helping the Brazilian Consumer Make a Healthier Choice. Food Sci Nutr Res. 2021; 4(2): 1-14.

## ABSTRACT

*We tested, in an online survey, how well five different front-of-pack (FOP) labels helped Brazilian consumers make a healthier choice between two food products as compared to a no FOP label control. All 1072 respondents were randomly allocated to one of six groups 1) no FOP label (control), 2) ABIA label, 3) GGALii Nutrient Profile label, 4) IdeC label, 5) Hybrid label or 6) Nutri-Score label and were all shown 9 food stimuli consisting of two products. The nutrient profiles of ABIA and Hybrid labels consider serving size of the food whereas the other three labels score per 100g. Respondents were asked which of the two products they perceived to be the healthier choice. Overall, the Hybrid and ABIA labels performed best, resulting in a statistically significantly higher percentage of correct answers as compared to the control for 9/9 and 8/9 of the food stimuli, respectively. Nutri-Score performed reasonably well and outperformed the control in 6/9 cases. The IdeC and GGALii NP warning labels were least helpful, outperforming the control group only once and twice, respectively. In conclusion, the Hybrid and the ABIA FOP labels, two interpretative traffic light labels that use colours and provide nutritional information per serving, were best suited to help Brazilian consumers choose the healthier product. They especially outperformed the other FOP labels when serving sizes differed significantly or when deeper consideration of nutritional information was needed to make an informed decision.*

## Keywords

Online consumer research, Front-of-pack label, Nutri-Score, Warning label, Traffic light label, Brazil.

## Introduction

Although many countries and health organizations have issued dietary guidelines with clear recommendations as to which food groups fit into a healthy diet and which foods and nutrients should be consumed in moderation, adherence to dietary recommendations is low.

Nutrient declarations on the back of the pack provide important information about the nutritional composition of food products, but consumers find this information complex and difficult to understand [1,2]. That's why, Front-of-Pack (FOP) nutrition labelling systems are designed to help the consumer make a quick, informed decision about the nutritional content or relative

healthfulness of a food or beverage.

There is no international standardization of FOP labels and therefore many different FOP labels have been developed and implemented across the world [3-5]. These FOP labels differ not only in graphic presentation, but also in the type and level of information they provide, the nutritional profiles used, the food products covered and the implementation (i.e. voluntary or mandatory). This is at least partly explained by the fact that the model must meet the needs for each country, taking into consideration cultural and dietary patterns, education level and alignment with national dietary guidance.

FOP labels can be roughly divided into two types namely interpretative and non-interpretative labels. Non-interpretative FOP labels provide information on the amount of key nutrients

(e.g. fat, sugar and sodium) and percent recommended intakes, with little interpretation of this information (e.g. Guideline Daily Amounts [GDA] or Reference Intake [RI]) logos [6]). Interpretative FOP labels contain an interpretation of the nutritional quality of the products, with or without information about the amount of key nutrients. Some FOP labels use colours (e.g. Multi Traffic Light [MTL] [7]) to indicate whether the level of a nutrient is high (red), medium (amber) or low (green). Other interpretive FOP labels provide a simple summary score of a product's overall nutritional profile (e.g. Keyhole logo [8], Choices Programme Logo [9]), a ranking (e.g. Health Star Rating [10], Nutri-Score [11]) or warn for high levels of certain critical nutrients in products (e.g. Warning labels [12]).

Many studies have examined the different attributes of effectiveness of FOP labels, but there are numerous inconsistencies in the results. This can be explained by the fact that the definition of effectiveness differs from study to study. Some studies focus on consumer liking, understanding or preference for FOP labels, others on food choice or actual or intended food purchase and some on food intake. There are also major differences in methodologies used. However, research into the effectiveness of FOP labels on consumer behavior in practice is lacking [13].

Studies focusing on consumers' understanding of FOP labels and product choice generally show that FOP labels appear to help consumers determine which foods are healthier and which are less healthy [14,15]. Simple FOP labels such as MTL, warning labels and Nutri-Score appear to be most effective [15].

The MTL label has been implemented in the United Kingdom [7]. It provides information on energy, fat, saturated fat, sugar and salt content per 100 gram and as percentage of reference intake, combined with traffic lights colours (green, amber and red) to highlight low, medium or high levels of the nutrients. It also provides serving size information that is expressed in easily recognizable and meaningful ways to the consumer (e.g.  $\frac{1}{4}$  of a pie). Nutri-Score is an interpretive FOP label that uses letters and colours to rank healthiness of products [11]. The French government adopted the Nutri-Score in 2017 and since then governments of other countries such as Belgium, Switzerland and Germany have also chosen to adopt the Nutri-Score [16,17]. In Latin America, the implementation of warning labels is spreading. Ecuador was the first to implement a mandatory FOP label system, i.e. a traffic light system [18]. They were followed by Chile which implemented mandatory warning labels in 2016 [19]. Since then, Peru [20], Paraguay and Uruguay also decided to implement warning labels and Mexico has recently followed suit [21]. Brazil recently reviewed mandatory nutrition labelling. Anvisa, the National Health Surveillance Agency of Brazil established a working group on Nutrition Labelling to identify problems in the transmission of nutritional information and alternatives that could help improve the effectiveness of nutrition labelling [22]. Several FOP labels were proposed to Anvisa and reviewed. In October 2020, ANVISA approved a FOP label for food and beverages, which is a warning label that uses a nutrient profile based on added

sugar, saturated fat and sodium content per 100g or 100 ml of product [23,24].

This study was performed in 2019 and designed to determine which of five different types of nutritional front-of-pack labels best helps Brazilian respondents identify the healthiest choice between two food products, compared to a non-label control. Four of the tested FOP labels were also reviewed by Anvisa. We wanted to test the robustness of the efficacy of different FOP labels by also comparing products from different product categories, consumed in different serving sizes or with closer nutritional profiles. As a result, wide range of products were tested.

## Materials and Methods

### Study population

Study participants were recruited from an existing research panel of Brazilian consumers that represent general members of the public. The aim was to have a representative sample of respondents and an even gender split. Participants were eligible for participation if they were aged between 18 and 65 years.

### Front-of-pack labels

Five FOP labels were tested in this study, i.e. GGALI<sup>ii</sup> Nutrient Profile, IDEC, ABIA, Nutri-Score and a hybrid label which was developed for this study. The first four FOP labels were selected because they were proposed by different stakeholders to Anvisa, the National Health Surveillance Agency of Brazil [22]. The selected labels cover three different visual expressions of nutritional labelling. GGALI<sup>ii</sup> Nutrient Profile and IDEC labels are warning style labels, Nutri-Score is an interpretative color coded label. The ABIA label is a traffic light system that provides nutritional information. The hybrid label is an adaptation of the Evolved Nutrition Label [25] and contains both nutritional information and a red color code, as warning sign, if levels of an ingredient are high. The Control Group received visual expressions of products without a FOP label. This group was used as a reference.

This study focused on the three nutrients of concern that were initially proposed by Anvisa, i.e. saturated fat, sugar and sodium [22]. Hence other nutrients were not taken into account.

### GGALI<sup>ii</sup> NP label

GGALI (Gerência-Geral de Alimentos) is Anvisa's General Food Management who prepared the "regulatory impact analysis" published by Anvisa in 2018 [22]. GGALI proposed two nutrient profiles. We selected the stricter one – GGALI<sup>ii</sup> –. The GGALI<sup>ii</sup> NP label is a warning style label highlighting high levels of nutrients of concern. It is based on the nutrient content per 100 g or 100 ml for food and beverages, having as reference the guidelines of the World Health Organization (WHO) and Codex Alimentarius [22]. Criteria were defined for low, medium and high content of free sugars, saturated fat, total fat and sodium [22]. For the GGALI<sup>ii</sup> NP label in this study criteria for high levels were applied. When this study was designed and conducted, Anvisa had not chosen the nutrient profile or visual model. The nutrient profile that was approved by ANVISA in 2020, is more lenient than the nutrient

profile that we used in this study. The magnifying glass visual that we used is similar to the visual approved by Anvisa [23,24]. This visual is also under discussion in Canada.

### IDEC label

IDEC (Instituto Brasileiro de Defesa do Consumidor) is a civil society in Brazil. They proposed a FOP label that is a warning style label and uses black triangles to inform the high content of sugars, total fat, saturated fat and sodium, and the presence of trans fats and sweeteners. The nutritional profile model was adapted from the Pan American Health Organization (PAHO) profile model and is based on percentage of energy [22].

### ABIA label

ABIA (Associação Brasileira das Indústrias da Alimentação), representing the Brazilian food industry sector, proposed a Multiple Traffic Light (MTL) FOP label, based on the United Kingdom traffic lights, which reports the absolute quantities of sugars, saturated fats and sodium per serving. It uses the red, amber and green colours to indicate the high, medium and low levels of each nutrient according to criteria per serving [22].

### Nutri-Score

Nutri-Score is an interpretative, graded, color-coded FOP label that has been developed by French researchers [11]. It is based on the nutrient profiling system of the United Kingdom Food Standards Agency which uses the nutrient content per 100 g for food and beverages. Positive points (0-10) are allocated for energy, total sugar, saturated fat and sodium content and negative points (0-5) are allocated for fruit, vegetables and nuts, fibre and protein content. Products scores range from -15 (most healthy) to +40 (least healthy) [11] and are translated into five categories of nutritional quality ranging from A (green) to E (red).

### Hybrid label

The hybrid FOP label is a model developed for this study and is

an adaptation of the Evolved Nutrition Label (ENL) [25]. Criteria for sugar, saturated fat and sodium were defined for small serving sizes (<60 g), medium serving sizes (60-120 g) and large serving sizes (>120 g). In line with ENL, calculations were done per serving size, except for serving sizes between 60 and 120 g, in which case calculations were done per 100 g. The label provides quantitative nutritional information per serving and uses the red color to indicate high amounts of the nutrient in a serving of the product.

An example of the five labels is provided in Figure 1. The specific criteria used for the ABIA, GGALI<sup>ii</sup>, IdeC and Hybrid labels are presented in Table S1. For the Nutri-Score label, the Nutri-Score algorithm was used to calculate the score for each product [11].

### Food stimuli

This study included 18 food items, which were presented to the respondents in sets of two. The respondents were asked to indicate which of the two products they thought was healthier. The correct answer was defined considering the contents of the three nutrients highlighted on the front of pack label. That is, the product with the lowest sugar, saturated fat and sodium content per serving was considered the healthiest choice. When a product was higher in one nutrient and lower in another, the larger difference was considered most important. In a few cases, one nutrient was slightly lower and one much higher. Subsequently the much higher nutrient was considered more important for the classification.

The food sets (food stimuli) were carefully selected to test the robustness of the labels to help the consumer identify the healthier option. The food stimuli differed with regard to the following variables: similar products consumed in small serving size, similar products consumed in large serving size, similar product but consumed in different serving sizes, and products from different food categories but consumed in same eating occasion. The food categories included in the study were soft cheeses, fat spreads, ice creams, lasagne, frozen meals, fermented milks & chocolate oat




Figure 1: Example of front-of-pack labels used in this study (translated from Portuguese to English).

drinks, sweet snacks, cereal bars & yoghurt and chocolate bars. The products corresponded to different consumption occasions (e.g. breakfast, lunch or main meal and in-between meal snack). Figure 2 shows an example of one of the stimuli as presented to the respondents. In Brazil it is not mandatory to declare sugar content of food products on the packaging. For some of the products used in this study, i.e. ice creams, frozen meals, the dairy alternative drink and sweet snacks, we had to estimate the sugar content. Estimations were based on similar products marketed in countries where sugar content is declared in the nutrition table. Table 1 summarizes the nutritional profiles of the food stimuli. Other, detailed information about the food stimuli can be found in Figure S1.

### Data collection

A 5-minute online survey was conducted in July 2019, using Toluna Quick Surveys. All respondents were randomly allocated to one of six groups 1) no label (control), 2) ABIA label, 3) GGALI<sup>ii</sup> NP label, 4) IDEC label, 5) Hybrid label or 6) Nutri-Score label.

Apart from the control group, the respondents were shown the FOP label. The FOP label was briefly explained. The respondents were then shown 9 food stimuli consisting of two products and, based on the information they received, they were asked which product they thought was the healthier choice. The respondents in the control group were also shown the same 9 choice sets of products, but without a label. Respondents were provided with four potential answers: 1) Product 1; 2) Product 2; 3) No difference and 4) I don't know. Respondents were also asked to rate, on a

scale from 0 to 10, how useful the label was in helping them make a healthy food choice and what they liked and disliked about the specific type of label that they had evaluated.

### Statistical analyses

Descriptive statistics were calculated for the sociodemographic data of the participants. The percentage of participants selecting the answers “Product 1”, “Product 2”, “No difference” or “Don't know” were calculated for each set of food stimuli, for each FOP label group as well as for the control group. Significance testing (Z-test) was performed to test if the proportion of participants correctly identifying the healthier product differed between the FOP label groups. Significance tests were performed within these subgroups to test whether participants who correctly chose the healthier option differed according to education level or income. The mean scores for usefulness of the labels were calculated. T-tests were used to test for statistical differences between mean scores.

Statistics were performed with the Toluna Analytics tool. A significance testing at a 95% confidence level was used.

## Results

### Respondents

A total of 1072 Brazilian men and women participated in the online survey. Sociodemographic data are presented in Table S2. A total of 176 respondents were included in the control group, 181 respondents were allocated to the ABIA group, 177 respondents



Figure 2: Example of stimulus used for each randomized group.



**Table 1:** Nutritional profile of food stimuli.

Food category	Product	Serving size (g)	Per serving			Per 100 g		
			Sugar (g)	Saturated fat (g)	Sodium (mg)	Sugar (g)	Saturated fat (g)	Sodium (mg)
Soft Cheese	Product 1*	30	NA	1,9	118	NA	6,3	393
Soft Cheese	Product 2	30	NA	5,2	239	NA	17,3	796
Fat spreads	Product 1*	10	NA	0,9	70	NA	9	700
Fat spreads	Product 2	10	NA	4,8	90	NA	48	900
Ice cream	Product 1	86	21,5	7,7	40	25	8,9	46,5
Ice cream	Product 2*	60	13	1,8	12	21,7	3	20
Lasagne	Product 1	400	12	10	1280	3	2,5	320
Lasagne	Product 2*	400	11,2	5,2	1440	2,8	1,3	360
Frozen meals	Product 1*	300	NA	2,9	250	NA	0,9	83,3
Frozen meals	Product 2	275	NA	9,3	1242	NA	3,4	451,6
Fermented milk drink	Product 1	200	32,5	0	75	16,25	0	37,5
Chocolate oat drink	Product 2*	260	12	0,6	60	4,6	0,2	23
Sweet snacks	Product 1	40	10,6	3,4	68	26,6	8,4	170
Sweet snacks	Product 2*	20	6,6	2,1	26	33	10,5	130
Cereal Bar	Product 1*	21	6,9	0,8	0	32,8	3,8	0
Yoghurt	Product 2	170	20,4	4,6	160	12	2,7	94
Chocolate bars	Product 1*	16,7	8	2,6	16	47,9	15,5	100
Chocolate bars	Product 2	40	19,9	6,6	40	49,8	16,5	100
* Healthier option								

**Table 2:** Percentage of participants correctly choosing the healthiest option within each food stimulus, by randomization assignment to FOP label or control.

	Soft cheese	Fat spreads	Ice cream	Lasagne	Frozen meals	Milk drinks	Sweet snacks	Cereal bar & yoghurt	Chocolate bars
Control	76 <sup>d</sup>	37 <sup>c</sup>	31 <sup>d</sup>	48 <sup>d,f</sup>	13	40	26 <sup>c</sup>	44 <sup>c,f</sup>	39 <sup>c,f</sup>
ABIA	88 <sup>a,c,d</sup>	82 <sup>a,c,d,f</sup>	85 <sup>a,c,d,f</sup>	46 <sup>d,f</sup>	81 <sup>a,c,d,f</sup>	73 <sup>a,c,d</sup>	77 <sup>a,c,d,f</sup>	85 <sup>a,c,d,f</sup>	76 <sup>a,c,d,f</sup>
GGALI <sup>ii</sup>	77 <sup>d</sup>	24	65 <sup>a,d</sup>	41 <sup>f</sup>	59 <sup>a,d</sup>	32	15	31	29
IdeC	51	32	13	33 <sup>f</sup>	12	34	23	70 <sup>a,c,f</sup>	33
Hybrid	88 <sup>a,c,d</sup>	75 <sup>a,c,d,f</sup>	88 <sup>a,c,d,f</sup>	64 <sup>a,b,c,d,f</sup>	85 <sup>a,c,d,f</sup>	78 <sup>a,c,d</sup>	79 <sup>a,c,d,f</sup>	86 <sup>a,c,d,f</sup>	72 <sup>a,c,d,f</sup>
Nutri-Score	87 <sup>a,c,d</sup>	58 <sup>a,c,d</sup>	60 <sup>a,d</sup>	20	67 <sup>a,d</sup>	74 <sup>a,c,d</sup>	66 <sup>a,c,d</sup>	22	26

Performing significantly ( $p < 0.05$ ) better than <sup>a</sup>: Control; <sup>b</sup>: ABIA; <sup>c</sup>: GGALI<sup>ii</sup> NP; <sup>d</sup>: IdeC; <sup>e</sup>: Hybrid; <sup>f</sup>: Nutri-Score within the same category.

were allocated to the GGALI<sup>ii</sup> NP warning label, 181 respondents to the IDEC – triangle warning label, 178 respondents to the hybrid color code label and 179 respondents were allocated to the Nutri-Score label.

There were no consistent significant differences in terms of education and income levels between the different groups.

### Accuracy of choosing the healthier product

Table 2 summarizes for each of the labels and the control group the percentages of respondents that correctly identified the healthier product for each of 9 stimuli.

When the soft cheese food stimuli were presented, most respondents accurately identified the healthier product. Respondents who were shown the IDEC warning label performed significantly worse than the control group; a third of the respondents in this group indicated that there was no difference between the two products and only 51% of the respondents correctly identified the healthier product.

Respondents who were shown the ABIA or hybrid labels were most likely to identify the healthier choice from the fat spreads category. Most respondents who were shown the GGALI<sup>ii</sup> or IDEC warning style labels selected the least healthy product of the

two (41% and 44%, respectively) or indicated that there was no difference between the two products (32% and 24%, respectively). The GGALI<sup>ii</sup> label scored worse than the control group. Also 55% of the respondents in the control group chose the less healthy product.

For the ice cream stimuli, the group of respondents who were shown the IDEC labels had most difficulty choosing the healthier option; only 13% of the respondents chose the healthier product. Forty percent of the respondents in the control group and 60% of the respondents from the IdeC group indicated that there was no difference between the two ice cream products. The ABIA and hybrid labels helped the respondents best to make the healthier choice, with 85% and 88%, respectively selecting the healthier product.

Many respondents had difficulty identifying the healthier lasagne. Only the respondents who were shown the hybrid label performed significantly better (64% correct) than the control group (48% correct). The Nutri-Score label helped only 20% of respondents make the right choice, compared to 48% of the respondents in the control group. Seventy percent of the respondents who were shown the Nutri-Score label indicated that there was no difference between the products.

---

Respondents in the control group and those shown the IDEC label found it particularly difficult to choose the healthier option from the frozen meals. A total of 56% of the respondents in the control group and 62% of respondents in the IDEC label group indicated that there was no difference between the two frozen meals. The ABIA and hybrid label performed best.

In the category fermented milk and chocolate oat drinks, the ABIA, Hybrid and Nutri-Score labels performed best and the GGALI<sup>ii</sup> and IDEC labels performed worse with about one-third of respondents choosing the less healthy option and about one-third indicating that there was no difference between products.

There was a marked difference between labels in their ability to help choose the healthier sweet snack. The majority of the respondents in the control group (56%) and those who were shown the GGALI<sup>ii</sup> (73%) or Idec (57%) label considered that there was no difference between the two products with regard to health. The labels ABIA, Hybrid and Nutri-Score performed significantly better.

When respondents were shown a cereal bar and a yoghurt product, i.e. two very different products, the ABIA, IDEC and hybrid labels were most successful in helping them make a healthier choice. Forty-five percent of respondents who were shown the GGALI<sup>ii</sup> label believed that there was no difference between products and 72% of respondents who were shown the Nutri-Score label selected the less healthy product.

There was confusion amongst the majority of the respondents for the chocolate bar category (showing two different sizes of the same brand chocolate bar). Most respondents seeing the GGALI<sup>ii</sup> (59%), Idec (56%) or Nutri-score (63%) labels, which do not consider serving size, thought that there was no difference between the two products. The ABIA and Hybrid labels, which do consider serving size, resulted in the highest numbers of respondents choosing the healthier option.

Overall, the IDEC warning label was least helpful for consumers to make the healthier choice. Eight out of nine times, the IDEC label for the healthier product was the same as for the less healthy product, providing no guidance to the consumer. Only once, when comparing the cereal bar and yoghurt, the IDEC label outperformed the control group. In that case, 70% of the respondents chose the healthier option. The other warning label, GGALI<sup>ii</sup> NP, performed a bit better than the IDEC label, but also failed to distinguish products six out of nine times. It outperformed the control group in only two out of nine cases. Nutri-Score performed reasonably well but also failed two times in guiding the consumer to the healthier choice when products were given the same rating. This happened for example when the serving sizes of the two products differed significantly. In case of the cereal bar (21g) versus yoghurt (170g) food stimulus, a better Nutri-Score was given for the least healthy product (yoghurt).

The hybrid label performed best, resulting in statistically significantly higher percentage of correct answers as compared to

the control in all cases. The ABIA label outperformed the control group eight times out of nine.

If we look at the participants who correctly identified the healthier options, there appears to be no effect of education level or income. Statistical tests showed a statistically significant effect for only 3 of the 36 subgroups tested (9 food stimuli x 6 FOP label groups). Low-income participants from the control group scored significantly higher than high-income participants in selecting the healthier ice cream. Participants with a higher income who were shown the ABIA label scored significantly higher than participants with a lower income in selecting the healthier milk drink. Finally, the less educated in the control group scored higher than the higher educated when selecting the healthier chocolate bar.

### Usefulness ratings

Respondents were asked to indicate on a scale of 0-10 how useful the label was in helping to choose the healthier product. Between 89% and 92% of respondents rated their label as either very useful (7-8) or extremely useful (9-10). Mean usefulness scores were 9.7, 9.6, 9.4, 9.6 and 9.8 for the ABIA, GGALI<sup>ii</sup> NP, IDEC, Hybrid and Nutri-Score labels, respectively, and did not differ statistically significant. The IDEC and GGALI<sup>ii</sup> NP warning labels had the highest proportion (7% and 5%, respectively) of respondents indicating that the label was not at all useful (0-4). The proportion of respondents that rated the IDEC labels as not at all useful (7%) was significantly higher than the 2% of respondents rating the Nutri-Score and Hybrid label as not useful. See Figure 3 for usefulness ratings of FOP labels.

### Feedback on labels

To better understand how the FOP labels were perceived by the respondents, they were all asked what they liked or disliked about the FOP label they were shown.

### ABIA label


Most respondents said they liked the label, especially the use of colors and the clarity and objectivity of the information provided. When asked about what they did not like, some respondents indicated that they would also like information about other nutrients and energy.

### GGALI<sup>ii</sup> NP label

Respondents were pleased with the label's simplicity, its decisive message and the fact that it attracts attention. Not all respondents were satisfied with the black color and some were missing nutritional information.

### Hybrid label

Respondents were particularly pleased with the use of the red color as it draws attention and also the clear and easy to understand information about the nutrients and serving size. However, the language should be kept simpler (e.g. salt instead of sodium) and some respondents wanted more information about other nutrients and energy.



**Answer to question:** To what extent is the label below useful when helping you make healthy food choices?

**Figure 3:** Usefulness ratings of the five FOP labels.

### IDEC

Respondents were especially pleased with the simple information that helps people make a quick decision. Not all respondents were happy with the black color and said it wasn't noticeable, and some would like more specific information about the amount of the nutrients in the products.

### Nutri-Score

When evaluating the Nutri-Score label, the respondents indicated that they liked the simple, clear message and the use of colors. Negative aspects of the Nutri-Score were the lack of information about nutrient levels and the underlying model. Some respondents misinterpreted the label as they believed that the colors and letters represented the presence of vitamins in the products.

In summary, respondents prefer simple FOP labels that use colors (not black) to convey the message. They would like to receive information about the amount of nutrients in the products, but in simple language.

### Discussion

FOP labels are designed to help consumers choose healthier food and drinks. This study compared how well five different FOP labels helped Brazilian consumers make a healthier choice between two food or drink products. It showed that the Hybrid label and the ABIA label, performed best under the tested conditions. Both labels provide both nutritional information per serving as well as concise interpretation using colors (i.e. traffic light or red light only).

Most of the other studies that investigated how well different FOP labels helped consumers make a healthier choice between

products, compared products within the same food category and with the same serving size, but with marked differences in nutrient profiles. These studies often found that simple interpretative labels such as Nutri-Score or warning labels were effective in helping the consumer make a healthier choice [26-29]. This makes sense, because when a label clearly distinguishes between products (e.g. different score, color or with or without a warning label), the consumer can easily make a choice. However, this does not reflect the complexity consumer's face when shopping as this clear distinction does not always exist. Our research showed that when labels do not clearly distinguish between two products, or when products from different product categories or with different serving sizes are compared, these simple labels do not help the consumers to make an informed choice.

Many factors influence how consumer's process information on a FOP label and how deeply this information is processed [30]. Consumers may only glance at the FOP label, process partial information or process the FOP label in depth. For example, the level of nutritional knowledge influences the type of information the consumer processes. Knowledgeable consumers are more likely to use the more complex nutrient information on complex labels, while a less knowledgeable consumer may look for calorie and color-coded information. Average consumers are more likely to process the information on the FOP label in depth. Under time pressure consumers will only quickly inspect the information on the FOP label and not process all available information [30]. Health-motivated consumers may also look more actively for nutritional information, while hedonically-motivated consumers may not look at nutritional information, but more at brand names [30]. So, depending on the situation, different types of FOP labels can be the most effective.

Our study showed that simple summary labels are effective when there is a clear distinction between products, but if a quick decision cannot be made, consumers will consider the nutritional information on the FOP label, when available. In those cases, interpretative labels that provide nutritional information to the consumer better assist the consumer in making an informed choice. This study showed that, even when the colors on the Hybrid and ABIA labels did not differ between products, participants could choose the healthier option based on the nutritional information presented on these labels. This suggests that the nutritional information facilitates the comparison of the nutritional content of the products, allowing the consumer to make a healthier choice. The two warning labels, i.e. IDEC and GGALI<sup>ii</sup> NP were in most cases not sensitive enough to help the consumer distinguish products based on healthiness. They did not outperform the control group. Both the Idec and GGALI<sup>ii</sup> labels use very strict nutrient profiles and therefore most products bear the logo [22], making them less sensitive to distinguish products. The nutrient profile that will be implemented in Brazil is more lenient and if we would have used this more lenient profile, the number of warning labels would have been different for two food stimuli; only one of two frozen meals would have carried a warning logo for sodium and the yogurt would have carried no warning label for added sugar. Besides the lack of discrimination between products, these labels are also very simplistic and do not contain additional nutritional information to help the consumer make an informed decision, where the number of warning labels for sugars, saturated fats and sodium does not differ between two foods.

When labels on two different products are the same, respondent interpret this as if products are equally healthy, or respondents base their decision on other information that they have about the product (e.g. packaging, type of product, claims on product, presence of other ingredients, and knowledge of the brand). For example, the vegetable-oil based spread used in this study contained less saturated fat and sodium than the presented butter, so it is nutritionally, the healthier choice. However, Brazilian Dietary guidelines promote butter consumption, not vegetable-oil based spreads. Butter can therefore be seen as more natural and healthier than vegetable-oil based spreads in Brazil and this is also reflected in the results. When the FOP labels indicated that the vegetable-oil based spread was the healthier choice, most respondents chose that product. However, if no label was shown, or if labels on butter and vegetable-oil based spreads did not differ (in case of IDEC and GGALI<sup>ii</sup> NP), more than 40% of participants chose butter as the healthier option.

A recent review of FOP schemes performed by the European Commission concluded that FOP schemes providing nutritional information per 100g were better understood than portion-based schemes [13]. However, more than 90% of the food categories in Brazil have regulated serving sizes less than 100g / 100ml. When a nutrient profile is standard applied in 100g or 100mL distorted comparisons are generated. For products consumed in serving sizes <100g or ml, the amount of nutrients to calculate the FOP label is overestimated, while for products consumed in portions

of >100g/ml it is underestimated. As a result, some products with small serving sizes will unfairly receive a warning label, while some products with large serving sizes that are high in nutrients of concern receive no warning label. For example, in this study, two lasagnas with a 400g serving size were compared. According to the nutritional profile criteria defined by GGALI<sup>ii</sup>, based on 100g, neither of the two products would receive a warning label, and with Nutri-Score both lasagnas would receive a score of B. These two FOP labels would thus suggest that products are healthy, despite the relatively high saturated fat and sodium contents per serving as % GDA.

While the GGALI<sup>ii</sup> NP and IDEC warning labels were the least successful in helping participants make the healthier food choice, the labels were considered by the respondents to be as useful as the other FOP labels. It is important to note that the respondents did not receive any feedback on how well they did. So, they were not aware of the correct answer and how often they correctly identified the healthier option or mistakenly assumed there was no difference. One could speculate that if they got this feedback, ratings of usefulness would be lower. In any case, the ratings show that any FOP label that could help make the consumer an informed choice is considered useful by consumers. For research purposes, asking this question without providing feedback to the participants does not seem relevant. Feedback from the participants suggests that simple FOP labels that use bright colors and contain nutritional information in simple language, are liked.

Grunert et al hypothesized that consumers' liking for FOP labels is guided by three considerations: 1) consumers like simplicity, 2) when provided with simplified information consumers still want to know what it stands for and how the simplified message (e.g. warning- or health logo) has been derived, and 3) nutrition information can create a consumer resistance when they feel pushed to make choices that they do not want to take [31].

This is also confirmed by a recent study conducted by Talati et al. [32] who investigated consumer perception of five FOP labels, i.e Health Star Rating, MTL, Nutri-Score, RI and a warning label. The colored FOP labels MTL and Nutri-Score stood out and were most liked by consumers in all countries. Although the most simplified FOP labels, Nutri-Score and warning labels, were easy to understand, they were perceived as providing insufficient information and the least trusted. The RI label was perceived as the most confusing but scored high on trust. Overall, the MTL label, which combines nutrient-specific information and a summary interpretation using colour, was most liked and trusted in this study.

A strength of the current study was that it really tested the robustness of five FOP labels that differed not only in visual expression and the amount of information provided, but also in the underlying nutrient profile. Unlike other studies that mostly tested products within the same food category and with the same serving size, this study was designed to compare how well these five FOP labels enabled consumers to choose between products



that differ in nutritional composition, serving size and/or food category as consumers face in real life. Another strength is that a control group has been included and that the effectiveness of the FOP labels in helping the consumer to choose the healthier product could therefore be compared with a reference group that was not given a FOP label. This provides insight into whether the presence of a specific FOP label is of added value for a consumer when making an informed choice.

Participants were also asked to select the healthier product. This demonstrates how effective the FOP label is in helping the consumer make a choice and whether it fits its purpose. In other studies consumers were asked which product they would buy [33-37], but this may be influenced by factors such as familiarity and liking of the product and cost of the product [31]. Other studies only asked which FOP label is preferred [32,38]. As demonstrated by our study, all FOP labels were rated as very useful, irrespective of their efficacy in helping the consumer choose the healthier option. Only asking for preference is thus not very useful.

This study also has some limitations. Participants were a representative sample of the Brazilian population. Therefore, we also included participants with a lower education level, who may have had difficulty understanding the information on the FOP labels. However, socio-economic status and level of education was similar between the six FOP label groups and therefore we did not expect this to affect the outcomes of the study. This was confirmed by statistical subgroup analyses showing that participants who correctly identified the healthier options, did not differ with respect to level of education or income. Another limitation is that we did not ask the participants if they were color-blind. Thus, it is possible that participants with color-blindness were included, which may have adversely affected the ability to understand the color-coded labels. However, none of the participants who were shown the ABIA or Nutri-Score labels, voluntarily reported being color-blind and thus unable to interpret the labels.

The FOP labels that we tested in this online survey were selected because they were under consideration by Anvisa, the National Health Surveillance Agency of Brazil, at the time we designed this study. Anvisa proposed in its preliminary report on the regulatory impact analysis on nutrition labelling [22] to focus only on the three nutrients of concern, sugar, saturated fat and sodium. We therefore decided to only use the content of these three nutrients to inform the different FOP labels (with exception of Nutri-Score). Focusing on just these three nutrients of concern is a limitation to assessing the healthiness of a product.

Brazilian regulation do not require the sugar content of food products to be stated on the packaging. For some of the products used in this study (ice creams, frozen meals, dairy alternative drink, sweet snacks), we had to estimate the sugar content. These estimates were unlikely to deviate very much from the actual sugar content and were used for all FOP labels.

This study was conducted online using pictures of actual

products. It does therefore not reflect a real-life situation in which participants can examine packaging and other information, such as the nutrition table on the back, to make an informed choice. Finkelstein et al [39] attempted to mimic a real-life situation by asking the participants (n=147) to purchase their weekly groceries in an online grocery store with 3343 foods and 832 beverages. Participants had only access to back-of-pack Nutrition Information Tables or were also shown an MTL label or Nutri-Score label. Both the MTL and Nutri-Score FOP labels improved the dietary quality of the purchases as compared to the control group. The Nutri-Score label performed best in improving overall diet quality, but unlike Nutri-Score, the MTL label reduced calories. Thus, FOP labels had added value when purchasing products, even in the presence of a Nutrition Information Table.

## Conclusions

In conclusion, this study showed that the Hybrid and the ABIA FOP labels, two interpretative labels that use colors and provide nutritional information per serving, were best suited to help Brazilian consumers choose the healthier product. The other three labels are based on per 100g and therefore will not always discriminate enough to help consumers. The ABIA and Hybrid labels outperformed the other FOP labels when serving sizes differed significantly or when deeper consideration of nutritional information was needed to make an informed decision.

## Acknowledgments

We thank the volunteers who participated in this online survey and all other staff at Harris Interactive, Toluna and Unilever who helped with the study or provided input to this manuscript.

## References

1. Cowburn G, Stockley L. Consumer understanding and use of nutrition labelling: a systematic review. *Public health nutrition*. 2005; 8: 21-28.
2. World Health Organization. Guiding principles and framework manual for front-of-pack labelling for promoting healthy diets. World Health Organization. Geneva. 2019.
3. Kanter R, Vanderlee L, Vandevijvere S. Front-of-package nutrition labelling policy: global progress and future directions. *Public health nutrition*. 2018; 21: 1399-1408.
4. Labonté M.È, Poon T, Gladanac B, et al. Nutrient Profile Models with Applications in Government-Led Nutrition Policies Aimed at Health Promotion and Non communicable Disease Prevention: A Systematic Review. *Advances in Nutrition*. 2018; 9: 741-788.
5. Van der Bend D.L.M, Lissner L. Differences and Similarities between Front-of-Pack Nutrition Labels in Europe: A Comparison of Functional and Visual Aspects. *Nutrients*. 2019; 11: 626.
6. The Food and Drink Federation. Food labelling.
7. Food Standards Agency. Front of Pack nutrition labelling guidance.

8. Swedish National Food Agency. The Keyhole.
9. Choices International Foundation. Choices programme.
10. Australian Government of Health. Front-of-pack labelling updates.
11. Julia C.R, Hercberg S. Development of a new front-of-pack nutrition label in France: the five-colour Nutri-Score. *Public Health Panorama*. 2017; 3: 712-725.
12. Reyes M, Garmendia M.L, Olivares S, et al. Development of the Chilean front-of-package food warning label. *BMC public health*. 2019; 19: 906.
13. Storksdiack genannt Bonsmann S.M.G, Ciolo E, van Bavel R, et al. Front-of-pack nutrition labelling schemes: a comprehensive review. Publications Office of the European Union. Luxembourg. 2020.
14. Cecchini M, Warin L. Impact of food labelling systems on food choices and eating behaviours: a systematic review and meta-analysis of randomized studies. *Obesity Reviews*. 2016; 17: 201-210.
15. Temple N.J. Front-of-package food labels: A narrative review. *Appetite*. 2020; 144: 104485.
16. Federal public service Health Food Chain Safety and Environment Belgium. Nutri-Score.
17. Bundesministerium für Ernährung und Landwirtschaft. Ergebnis der Verbraucherbeteiligung liegt vor: Bundesministerin Julia Klöckner wird Nutri-Score(R) einführen. 2020.
18. Cox Vásconez A.E. RESOLUCIÓN No. 14 511. Gobierno Nacional de la Republica del Ecuador. Ministerio de Industrias Y Productividad. Subsecretaria de la Calidad. Quito. Ecuador. 2014.
19. Carreño I. Chile's Black STOP Sign for Foods High in Fat, Salt or Sugar. *European Journal of Risk Regulation*. 2015; 6: 622-628.
20. El Presidente de la Republica Peru. Aprueban Manual de Advertencias Publicitarias en el marco de lo establecido en la Ley N° 30021, Ley de promoción de la alimentación saludable para niños, niñas y adolescentes, y su Reglamento aprobado por Decreto Supremo N° 017-2017-SA. *El Peruano*: Peru. 2018.
21. Rojo Sánchez A.G.N.B, J.A. Modificación a la Norma Oficial Mexicana NOM-051-SCFI/SSA1-2010, Especificaciones generales de etiquetado para alimentos y bebidas no alcohólicas preenvasados-Información comercial y sanitaria, publicada el 5 de abril de 2010. Secretaria De Economía Gobierno Mexicano, Ed. Mexico City 2020; Vol. NOM-051-SCFI/SSA1-2010.
22. Agência Nacional de Vigilância Sanitária (Anvisa). Relatório Preliminar de Análise de Impacto Regulatório sobre Rotulagem Nutricional. Gerência-Geral de Alimentos; Brasília. 2018.
23. Didier D. Resolução de diretoria colegiada-RDC n° 429, de 8 de outubro de 2020- ANVISA. 2020.
24. Colegiada M.d.S.A.N.d.V.S.D. Instrução normativa-in n° 75, de 8 de outubro de 2020. Estabelece os requisitos técnicos para declaração da rotulagem nutricional nos alimentos embalados. *Health, M.o., Ed. Diário oficial da união*. 2020; 195: 113.
25. Egnell M, Kesse-Guyot E, Galan P, et al. Impact of Front-of-Pack Nutrition Labels on Portion Size Selection: An Experimental Study in a French Cohort. *Nutrients*. 2018; 10: 1268.
26. Egnell M, Ducrot P, Touvier M, et al. Objective understanding of Nutri-Score Front-Of-Package nutrition label according to individual characteristics of subjects: Comparisons with other format labels. *PLoS one*. 2018; 13: e0202095.
27. Egnell M, Talati Z, Gombaud M, et al. Consumers' Responses to Front-of-Pack Nutrition Labelling: Results from a Sample from The Netherlands. *Nutrients*. 2019; 11: 1817.
28. Galan P, Egnell M, Salas-Salvado J, et al. Understanding of different front-of-package labels by the Spanish population: Results of a comparative study. *Endocrinologia, diabetes y nutricion*. 2020; 67: 122-129.
29. Arrua A, Machin L, Curutchet M.R, et al. Warnings as a directive front-of-pack nutrition labelling scheme: comparison with the Guideline Daily Amount and traffic-light systems. *Public health nutrition*. 2017; 20: 2308-2317.
30. Sanjari S.S, Jahn S, Boztug Y. Dual-process theory and consumer response to front-of-package nutrition label formats. *Nutrition reviews*. 2017; 75: 871-882.
31. Grunert K.G, Wills J.M. A review of European research on consumer response to nutrition information on food labels. *Journal of Public Health*. 2007; 15: 385-399.
32. Talati Z, Egnell M, Hercberg S, et al. Consumers' Perceptions of Five Front-of-Package Nutrition Labels: An Experimental Study Across 12 Countries. *Nutrients*. 2019; 11: 1934.
33. Talati Z, Egnell M, Hercberg S, et al. Food Choice Under Five Front-of-Package Nutrition Label Conditions: An Experimental Study Across 12 Countries. *American journal of public health*. 2019; 109: 1770-1775.
34. Mora-Garcia C.A, Tobar L.F, Young J.C. The Effect of Randomly Providing Nutri-Score Information on Actual Purchases in Colombia. *Nutrients*. 2019; 11: 491.
35. Billich N, Blake M.R, Backholer K, et al. The effect of sugar-sweetened beverage front-of-pack labels on drink selection, health knowledge and awareness: An online randomised controlled trial. *Appetite*. 2018; 128: 233-241.
36. Machin L, Arrua A, Gimenez A, et al. Can nutritional information modify purchase of ultra-processed products? Results from a simulated online shopping experiment. *Public health nutrition*. 2018; 21: 49-57.
37. Acton R.B, Jones A.C, Kirkpatrick S.I, et al. Taxes and front-of-package labels improve the healthiness of beverage and snack purchases: a randomized experimental marketplace. *The international journal of behavioral nutrition and physical activity*. 2019; 16: 46.
38. Dana L.M, Chapman K, Talati Z, et al. Consumers' Views on the Importance of Specific Front-of-Pack Nutrition Information: A Latent Profile Analysis. *Nutrients*. 2019; 11: 1158.
39. Finkelstein E.A, Ang F.J.L, Doble B, et al. A Randomized Controlled Trial Evaluating the Relative Effectiveness of the Multiple Traffic Light and Nutri-Score Front of Package Nutrition Labels. *Nutrients*. 2019; 11: 2236.

## Supplementary Materials

**Table S1:** Criteria used for the ABIA, GGALI<sup>ii</sup> NP, IdeC and Hybrid labels.

Label	Product criteria	Category	Total sugar (g)	Free sugars (g)	Saturated fat (g)	Sodium (mg)
ABIA label	Solids<100g and Liquids (ml)	Low	≤ 5,0		≤ 1,5	≤ 80
ABIA label	Solids<100g and Liquids (ml)	Medium	> 5,0 ≤ 13,5		> 1,5 ≤ 3,0	> 80 ≤ 300
ABIA label	Solids<100g and Liquids (ml)	High	> 13,5		> 3,0	> 300
ABIA label	Solids ≥ 100 g	Low	≤ 5,0		≤ 1,5	≤ 80
ABIA label	Solids ≥ 100 g	Medium	> 5,0 ≤ 27,0		> 1,5 ≤ 6,0	> 80 ≤ 600
ABIA label	Solids ≥ 100 g	High	> 27,0		> 6,0	> 600
GGALI <sup>ii</sup> NP label	Solid (100 g)	Warning label		≥ 10	≥ 4	≥ 400
GGALI <sup>ii</sup> NP label	Liquids (100 mL)	Warning label		≥ 5	≥ 2	≥ 200
IdeC label#	All products	Warning label		≥ 10%E	≥ 10%E	≥ 1 mg/kcal
Hybrid label	Solids < 60 g and Liquids	High	> 13,5 per serving		> 3,0 per serving	> 300 per serving
Hybrid label	Solids ≥ 60g ≤ 120g	High	> 22,5 per 100 g		> 5,0 per 100 g	> 500 per 100 g
Hybrid label	Solids > 120 g	High	> 27 per serving		>6,0 per serving	> 600 per serving


# %energy of product

**Table S2:** Sociodemographic data


Characteristics	Number of participants (%)
<b>Sex</b>	
Female	578 (54%)
Male	494 (46%)
<b>Age</b>	
18-34	681 (64%)
35-54	338 (32%)
55+	53 (5%)
<b>Education</b>	
Low	82 (8%)
Medium	481 (45%)
High	496 (46%)
Undisclosed	13 (1%)
<b>Social grade</b>	
A	209 (19%)
B1	194 (18%)
B2	329 (31%)
C1	179 (17%)
C2	103 (10%)
D-E	22 (2%)
Undisclosed	36 (3%)
<b>Income</b>	
Low	596 (56%)
High	401 (37%)
Undisclosed	75 (7%)
<b>Geographic location</b>	
Midwest	60 (6%)
Northeast	227 (21%)
North	41 (4%)
South	145 (14%)
Southeast	581 (54%)
Undisclosed	18 (2%)
<b>Household shopping responsibility</b>	
Sole	54%
Joint	38%
None	8%

Figure S1: Overview of all stimuli.


**Food stimulus: Soft cheese**




**Food stimulus: Fat Spreads**



**Food stimulus: Ice-Cream**




**Food stimulus: Lasagna**






**Food stimulus: Frozen meals**

Escondidinho de carne      Strogonoff de frango




**Food stimulus: Fermented milk & chocolate oat drink**

Leite Fermentado      Bebida aveia e cacau




**Food stimulus: Sweet snacks**

Bolinho de morango      Cookie chocolate



**Food stimulus: Cereal Bar & Yoghurt**

Barra de cereal de chocolate      Iogurte com geleia de morango



## Food stimulus: Chocolate bars

